

Transmutation of Elements through Capture of Electrons by Nuclei

A. Mukherji^{*†}

Formerly of Saha Institute of Nuclear Physics,
Block AF, BidhanNagar, Kolkata, 700064, India

PACS no. 01.55.+b

Abstract

A proton can capture an electron and turn into a neutron provided the electron has a kinetic energy of 0.782 MeV or more. An element of the Periodic Table can change into another on being exposed to such high energy electrons.

1 Introduction

The possibility of converting Mercury into Gold intrigued scientists for ages. The alchemists made considerable efforts towards this end without success.

Transmutation of one element into another takes place in radioactive decays and in nuclear reactions involving emission or absorption of charged particles. Owing to poor yields these processes are not suitable for bulk production of an element at a commercially viable cost.

Atoms split and unite in nuclear fission and fusion. Such privileges are enjoyed exclusively by a few members of the periodic table at its two extremities.

^{*}Mailing address: D 111 Shanti Niket, 32/4 Sahitya Parishad Street, Kolkata, 700006, India

[†]Electronic address: a.mukherji@saha.ac.in

2 *K*-Capture

The nucleus of a heavy element can capture[1] a *K*-shell (1s) electron and thereby convert itself into a different element. This phenomenon is viewed as follows.

The attraction between a nucleus and its nearest neighbour 1s electron increases with increasing atomic number Z . The atomic orbital shrinks in diameter. The electron itself comes closer to the nucleus causing further increase in their mutual attraction. The net effect is that the force between them varies as Z^4 . When Z is large the attraction becomes strong enough to release the electron from its bondage to its orbital. It is then captured by the nucleus.

In this description the 1s electron is depicted as revolving around the nucleus following a specified path and maintaining a definite distance. This is, of course, far from being literally true. Every *s*-electron in an atom occasionally passes through the nucleus. Its wavefunction is non-zero at the location of the nucleus. Hence, it has a finite probability of being within the nuclear volume. This causes hyperfine structures in atomic spectroscopy through Fermi[2] δ -interaction. The nucleus only needs to hold on to a passing electron. Of all the *s*-electrons the 1s-electron is the most energetic one. It travels the fastest through the nucleus. It is reasonable to expect that the 1s-electron will be the most difficult one to catch. But in reality only *K*-shell electrons are captured by nuclei. The slower *s*-electrons are ignored. The disparity between logical expectation and physical observation calls for a second look into the matter.

The phenomenon of *K*-capture is discontinuous with respect to Z , while the electron-nucleus force is a continuous function of Z . Explanation of a discontinuous feature in terms of a continuous function is, in principle, liable to be faulty.

The secondary cosmic rays that reach earth's surface are mostly slow electrons. They are perpetually present in great abundance. They are free electrons and are not fettered to any orbital. They are easy prey for capture by any nucleus — heavy or light — on land or sea. No nucleus has ever been known to have captured a cosmic ray electron. Otherwise, dire consequences would have followed. Hydrogen nuclei everywhere on earth would have turned into neutrons and the bound electrons would have gone free. A great plasma of free neutrons and electrons would have resulted in stead of living beings.

3 Induced Transmutation

When a nucleus captures an electron one of its protons turns into a neutron. This change is permissible from the point of view of charge conservation but not of mass conservation. The sum-total of the masses of a proton (938.272 MeV) and an electron (0.511 MeV) falls short of the mass of a neutron (939.565 MeV) by 0.782 MeV. Hence proton to neutron conversion is prohibited unless the captured electron provides this extra mass in the form of kinetic energy. It is very large for an electron with rest mass of about 0.5 MeV and is in the relativistic range. No nucleus can, therefore, capture a slow electron. Only 1s-electrons in very heavy atoms can provide the required energy. Hence, K -capture is limited to them only. The minimum energy requirement is, however, subject to a small correction due to changes in binding energies of the nucleus and of the electrons in various orbitals associated with the change in the constitution of the nucleus.

Transmutation of an element X into another Y can be induced by irradiating X by a strong beam of high energy electrons of at least 0.782 MeV. The new element Y can be extracted from the exposed target by chemical means. The process is complete if the isotope of Y so produced is a stable one. Otherwise Y may undergo radioactive decay and get converted into some other element.

It is now a matter of technology and experimentation to produce any desired element with high yield at low cost. The output will be chemically indistinguishable from that obtained from natural sources. The difference will only be in their isotopic constituents.

If large quantities of gold and/or other noble metals could be produced cheaply, they would lose their elevated positions as asset materials. World economy will suffer a jolt and it will be in need of a new standard.

References

- [1] John M Blatt and Victor F Weisskopf, *Theoretical Nuclear Physics*, (John Wiley, New York, 1952)
- [2] C.P.Slichter, *Principles of Magnetic Resonance*, (Springer, Berlin, 1980)